



## Two general scenarios make us believe to consider clusters of galaxies as potential gamma-ray emitters:

## (1) Multifrequency evidence of nonthermal activity



### X-ray emission is IC of the radio emission producing electron by the CMB

- · power-law with index simply related to observed radio (synchrotron) emission
- · matching spatial emission profiles in X-ray- and radio (halo) images
- · multifrequency models require a distinct relativistic particle population





## (2) Large-scale cosmological structure formation

### Hierarchical merging scenarios of galaxy clusters:

Larger structures evolve from mergers of adjacent but \$maller structures

baryonic matter condenses in form of galaxy cluster's

dark matter halos interact/merge

particles at the shock fronts

1st order Fermi acceleration of (Starting z=2.5) (from Jenkins et al. 1998)



### Three ways to detect galaxy clusters at gamma-rays

Direct detection of gamma-ray emission in unambiguous positional coincidence with an candidate galaxy cluster -> "pointing" approach

Spatial-statistical correlation between still unidentified gamma-ray sources and a candidate object population -> "correlation study"

Unresolved gamma-ray emission through localized enhancements in the extragalactic background

-> "background contribution studies"



## @ diffuse day:

## Special emphasis on the diffuse fore- and background involved in galaxy cluster analysis

### -> NO discussion of individual cluster observations

```
(McGlynn, Vestrand & Jennings: 22 Abell clusters -> u.l.)
(Sreekumar et al. 1996: Virgo, Coma -> u.l.)
(Reimer et al. 2003: 58 nearby X-ray bright Abell clusters -> u.l.)
```

## -> NO dicussion of spatial-statistical correlation work

```
(Kolatt & Piran 1996: Abell clusters <-> GRBs)

(Marani et al. 1997: rich, nearby Abell clusters <-> GRBs)

(Hurley et al. 1997: Abell clusters <-> GRBs)

(Gorosabel & Castro-Tirado 1997: Abell <-> GRBs)

(Burenin, Sunyaev et al. 1997: Abell <-> GRBs)

(Hurley et al. 1999: Abell clusters <-> GRBs)

(Colafranceso 2001,2002: Abell clusters <-> EGRET unIDs)

(Kawasaki & Totani 2002: possible merging clusters <-> EGRET unIDs)

(Reimer et al. 2003: nearby X-ray bright Abell <-> EGRET unIDs)
```



# Galaxy clusters as explanation of the extragalactic diffuse gamma-ray background (EGDB)

... a long record as well:

Strong & Bignami 1983: NGC 1275 & Perseus cluster possible sources of

γ-ray enhancement measured by COS-B

and subsequently:

Houston et al. 1984: cosmic ray interactions with intergalactic gas within


groups and clusters of galaxies produced EGDB

but: respective COS-B fluxes not confirmed by CGRO

Dar & Shaviv 1995: Cosmic Ray Origin of EGDB, essentially 100% (!)

Coma, Perseus, Virgo  $\sim 10^{-7}$  cm<sup>-2</sup> s<sup>-1</sup> (!)





Berezinsky, Blasi & Ptuskin 1997

#### Predictions of the contribution from clusters to the EGDB

-> not detectable by

current instrumentation EGHET Colafrancesco & Rlasi 1998 -ES GLOGE BRIV CITY S' 1 St" - - - Unresolved point sources - - - - -<- Waxman & Loeb 2000 vF\_[keV/cm²/sec/sr] IR Abs. off 10-6 <- Totani & Kitayama 2000 Photon Energy [GeV] GLAST Collaboration Meeting, Sep 03



#### BULK SOURCE OF UNIVERSE'S GAMMA RAYS IDENTIFIED, SCIENTISTS SAY

Scientists at Columbia University and Barnard College have found that the majority of the gamma rays outside of our galaxy are likely emitted by galaxy clusters and other massive structures. This may resolve a 30-year-old mystery as to the origin of the Universe's gamma-ray background. ... "This result not only resolves the question of where all these gamma rays are coming from, but provides a new probe of the gravity-driven picture of structure formation in the Universe"...

Barnand College aubi

Actrophysics Journal sine-year second of gi space, collected by divoughout the 1990s

#### Statistical Detection of Galaxy clusters in EGRET data -> unresolved sources -> EGDB

Method: annular binning of EGRET intensity data

|b|> 45°, 2.7° cut around 100 identified EGRET sources

sample: positions of 2469 Abells, 447 Abell ( $R \ge 2$ ), stacked out to 20°

average indivit

What does this mean?

sample size: |b|> 45°: 3.68 sr, - 0.69 sr source contribution: ~3 sr

 $1/2 \pi r^2 ... \pi r^2$ :472 ...945 sr or 150 ... 300 x oversampling of a bin in EGRET intensity map

residuals in EGRET data:

sources below 3EG catalog detection threshold(s) exist

- + contributions of unidentified AGN (unIDs awaiting their AGN identification)
- + contributions of identified sources with psf-extension of > 2.7° (the psf tails!),


+ plain diffuse galactic foreground (unadapted GALDEF!)
GLAST Collaboration Meeting, Sep 03



Reimer et al. 1999 (ICRC '99), 2001 (y2001), 2003 (ApJ 588)

58 of the X-ray brightest clusters, nearby (z < 0.14) as well as in  $\frac{\text{superposition}}{\text{superposition}}$ 


naturally included := best-observed clusters: EUV excess; hard X-ray emission; most of the radio halo clusters; Perseus, Coma, Virgo






The superposition: a highly

-> stacking of data of all in (~ 650 indiv. vp's) in clus



- -> usage of an appropriate ( model for exactly \*this\* sample
- -> correction for different foreground expectations to account for unequal exposures  $c_i = \frac{1}{\varepsilon_{\text{tot}}} \sum_i \varepsilon_i \text{DF}_i$ ,  $\varepsilon_{\text{tot}} = \sum_i \varepsilon_i$ .
- -> max Ih algorithm
- -> flux determination at image center






## Galactic diffuse model as used in likelihood application (Hunter et al. 1997)

here incl. positions of considered clusters; E > 100 MeV; gmult = 1, gbias = 0




### the complete sample (58)

the reduced sample (50)
(predominant identified point sources or galactic plane at image center -> removed!)







### NO DETECTION!


combined exposure: 3.5 x 10<sup>10</sup> cm<sup>2</sup> s

upper limit (50 cluster sample): 5.9 x 10<sup>-9</sup> cm<sup>-2</sup> s<sup>-1</sup>



energy averaged psf (E > 100 MeV) 6 wrong ire 68% in 3.1° aperture, 1° corresponds only 24% flux enclosure

claim recalc using correct psf conclusion  $r=1^{\circ}$ :  $1.14 \times 10^{-9}$  cm<sup>-2</sup>s<sup>-1</sup>  $4.7 \times 10^{-9}$  cm<sup>-2</sup>s<sup>-1</sup> marginally inconsistent  $r=2^{\circ}$ ?  $9.6 \times 10^{-9}$  cm<sup>-2</sup>s<sup>-1</sup> our u.l.:  $5.9 \times 10^{-9}$  cm<sup>-2</sup>s<sup>-1</sup> clearly inconsistent !!



main result figure from S&M 2002



### Conclusions:

No observational evidence for the contribution of galaxy clusters at high-energy gamma-rays yet.

Analysis requires precise handling of diffuse foreground, instrumental response, observational pecularities, and statistical assessments (noise expectations) ... but EGRET was \*just\* not sensitive enough here

[as for radio galaxies & Seyferts (Cillis, Hartman & Bertsch 2003, in submission), Starburst galaxies (Blom et al. 1999) and Normal Galaxies (Pavlidou & Fields 2001)]



### Conclusions 2:

Confirmation of this negative observational result from most recent publications:

synthetic spectra for a Conna-type


Miniati 2002, 2003:

[20...30% EGDB, but accounted for p & e]

Keshet et al. 2003 [~ 10% EGDB]

Gabici & Blasi 2003 [< 10% EGDB, following cluster merger evolution]

Berrington & Dermer 2003 [Only minor contribution to EGDB, which is featureless power law of 2.1]  $_{\rm GLAST\ Collaboration\ Meeting,\ Sep\ 03}$ 






### Conclusions 3:

When a cluster will be detected at gamma-rays (by GLAST), predictions on the EGDB contribution will be more precisly determined/verified.

We were presumable close to the required instrumental sensitivity in EGRET, so it's a <u>very appropriate</u> science case for GLAST.

Predictions exist already! (see literature)





-> improved modelling of high-latitude  $\gamma$ -rays based on IC halo emission Stay tuned!

